

**DESCRIPTION:** 

Ciprofloxacin belongs to the group of quinolones and acts bactericidally against mainly Gramnegative bacteria like Campylobacter, E. coli, Haemophilus, Mycoplasma, Pasteurella and Salmonella spp.

# **INDICATIONS:**

Gastrointestinal, respiratory and urinary tract infections caused by ciprofloxacin sensitive microorganisms like Campylobacter, E. coli, Haemophilus, Mycoplasma, Pasteurella and Salmonella spp. in calves, goats, poultry, sheep and swine.

# **CONTRA INDICATIONS:**

Hypersensitivity to ciprofloxacin.

Administration to animals with a serious impaired hepatic and/or renal function.

Concurrent administration of tetracyclines, chloramphenicol, macrolides and lincosamides.

# **SIDE EFFECTS:**

Hypersensitivity reactions.

Administration to juvenile animals can lead to arthropathy.

# WITHDRAWAL TIMES:

For meat: 12 days.

# **PACKAGING:**

Bottle of 100 ml, 500 ml, 1000 ml and 5000 ml

# DOSAGE:

For oral administration:

Poultry:

100 ml per 300 - 400 L of drinking water for 3 - 5 days.

Calves, goats and sheep:

Twice daily 5 ml per 75 - 150 kg body weight for 3 - 5 days.

A product of:





LG-113, Xth Central Mall, Mahavir Nagar, Kandivali (W), Mumbai 400067, Maharashtra, (INDIA). Ph: +91 22 3513 1576 / 77, Email : info@biochempharma.in, Web: www.biochempharma.in