Cobac Granules for oral administration

DESCRIPTION:

Colistin is an Poultry Antibiotics from the group of polymyxins with a bactericidal action against Gram-negative bacteria like *E. coli, Haemophilus* and *Salmonella*. Since Colistin is absorbed for a very small part after oral administration only gastrointestinal indications are relevant.

COMPOSITION:

Each gram contains:	
Colistin Sulphate	
(Equivalent to Colistin base)	4 800 000 I.U
Carrier	0.S

INDICATION:

Gastrointestinal infections caused by colistin sensitive bacteria, like *E.coli, Haemophilus and Salmonella spp.* in calves, goats, poultry, sheep and swine.

DOSAGE & ADMINISTRATION:

For oral administration : Calf, Goat and Sheep : Twice daily 1 g per 80 kg body weight for 5-7 days. Poultry and Swine : 100 g per 400-800 L of drinking water or 200-400 kg feed for 5-7 days

Note: For pre-ruminant calves, lambs.

WITHDRAWAL PERIOD:

For Meat : Minimum 28 days For Eggs : Minimum 7 Days

PRESENTATION:

100 g, 500 g & 1 Kg



LG-113, Xth Central Mall, Mahavir Nagar, Kandivali (W) Mumbai 400067, Maharashtra, (INDIA), Ph: +91-22-3513 1576/77 Email: info@biochempharma.in, Web: www.biochempharma.in