

# LEVOX-L

Levofloxacin 10% Oral Liquid

## COMPOSITION :-

Each ml contains :

Levofloxacin Hemihydrate USP

equivalent to levofloxacin.....100 mg

Approved colours.

Excipients.....Q.S.

## INDICATIONS:-

It is effective against both gram positive and gram negative bacteria, Mycoplasma and anaerobic organisms. Its wide spectrum of activity includes the pathogenic organisms like *E coli*, *Salmonella spp.*, *Staphylococcus spp.*, *Streptococcus spp.*, *Kiebsiells spp.*, *Pseudomonas aeroginosa*, *Bordetella bronchiseptic*, *Mycoplasma spp.*, *Clostridium perfringes* and *Camphylobacter*. Diseases produced by these organism in poultry can effectively be treated with Levofloxacin 10% Oral Liquid.

## DOSAGE & ADMINISTRATION:-

It is administered through drinking water at the following dosage rate -

**For treatment** : 1 ml per 1-2 L of drinking water or 20 mg/kg body weight for 3-5 days

**For Prevention**: 1 ml per 2-4 L of drinking water or 10 mg/kg body weight for 3-5 days

Or as directed by the registered veterinarian.

## TARGET SPECIES:-

Poultry

## WARNINGS:-

- Determine the body weight accurately before treatment, to ensure correct dosage.
- Wash the hands thoroughly after use.
- After treatment, animals should be moved to clean place in order to prevent re-infection.
- After the end of treatment, the watering system should be cleaned appropriately to prevent drug resistance.
- It should not be used for poultry resistant to any Quinolone derivative because cross resistance may develop.

## WITHDRAWAL:-

Meat: 4 days after the latest treatment prior to slaughter.

Egg: 7 days after the latest treatment prior to laying period.

## STORAGE:-

Store below 30° C. Protect from light.

A product of:

**Biochem** Pharma  
All we need is a healthy world



**BIOCHEM PHARMA**

LG-113, Xth Central Mall, Mahavir Nagar, Kandivali (W),  
Mumbai 400067, Maharashtra, (INDIA). Ph: +91 22 3513 1576 / 77,  
Email : info@biochempharma.in, Web: www.biochempharma.in

