Oxytetracycline 20%



Composition:

| Each Kg contains | |
|-----------------------------------|--------------|
| Oxytetracycline | 200 g |
| (equivalent to oxytetracycline hy | drochloride) |
| Carrier | Q.S. |

Benefits:

- OXYTRYN-200 is a broad-spectrum anti-infective that has been proven effective against a wide variety of infectious diseases caused by susceptible gram-positive and gram-negative bacteria.
- It is safe, stable and highly effective.
- It works in both the bloodstream and in the gastrointestinal tract.

Usage & Administration:

Increased rate of weight gain and improved feed efficiency: 10-50 g/ton feed continuously Control of infectious synovitis caused by *Mycoplasma synoviae*; control of fowl cholera caused by *Pasteurella multocida*: 100-200 g/ton (Feed continuously for 7-14 days)

Control of chronic respiratory disease (CRD) and air sac infection caused by *Mycoplasma gallisepticum* and *Escherichia coli*: 400 g/ton (Feed continuously for 7-14 days)

Reduction of mortality due to air sacculitis (air sac infection) caused by *Escherichia coli*: 500 g/ton (Feed continuously for 5 days).

WARNING:

At 500 g/ton level, withdraw 24 hours before slaughter.

Zero-day withdrawal period for lower use levels.

In low calcium feeds withdraw 3 days before slaughter.

Do not administer to chickens producing eggs for human consumption.

For use in dry poultry feeds only.



Bochem

A product of:

